

Neo-Oxy 100/100 MR[®]

..... NEOMYCIN SULFATE - OXYTETRACYCLINE HYDROCHLORIDE

Type A Medicated Article

FOR USE IN THE MANUFACTURE OF MEDICATED FEEDS

ACTIVE DRUG INGREDIENTS:

Oxytetracycline Hydrochloride 100 g/lb.
Neomycin Sulfate 100 g/lb.

MIXING AND USE DIRECTIONS

Mix Neo-Oxy 100/100 MR[®] with nonmedicated milk replacer to provide the following concentrations:

Indications for use	Oxytetracycline and Neomycin Amount	lb. of Neo-Oxy 100/100 [®] per ton of Type C Medicated Feed ¹
CALVES (milk replacer)		
For calves (up to 250 lb) for increased rate of weight gain and improved feed efficiency	0.05-0.1 mg/lb of body weight daily Feed continuously; in milk replacers or starter feed.	0.1-0.2 ²
RESIDUE WARNING: Zero day withdrawal. Use of more than one product containing neomycin or failure to follow withdrawal times may result in illegal drug residues.		
For calves (up to 250 lb) for treatment of bacterial enteritis caused by <i>E. coli</i> susceptible to oxytetracycline; treatment and control of colibacillosis (bacterial enteritis) caused by <i>E. coli</i> susceptible to neomycin	10 mg/lb of body weight daily Feed continuously for 7-14 days in milk replacers or starter feed. If symptoms persist after using 2 or 3 days, consult a veterinarian. Treatment should continue 24 to 48 hours beyond remission of disease symptoms.	20 ²
RESIDUE WARNING: A withdrawal period has not been established for use in preruminating calves. Do not use in calves to be processed for veal. A milk discard time has not been established for use in lactating dairy cattle. Do not use in female dairy cattle 20 months or older, including dry dairy cows. Use in these cattle may cause drug residues in milk and/or in calves born to these cows. Withdraw 5 days before slaughter. Use of more than one product containing neomycin or failure to follow withdrawal times may result in illegal drug residues.		
CALVES, BEEF CATTLE, AND NONLACTATING DAIRY CATTLE		
For treatment of bacterial enteritis caused by <i>E. coli</i> and bacterial pneumonia (shipping fever complex) caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline; treatment and control of colibacillosis (bacterial enteritis) caused by <i>E. coli</i> susceptible to neomycin.	10 mg/lb of body weight daily Feed continuously for 7-14 days in feed or milk replacers. If symptoms persist after using 2 or 3 days, consult a veterinarian. Treatment should continue 24 to 48 hours beyond remission of disease symptoms.	20 ²
RESIDUE WARNING: A withdrawal period has not been established for use in preruminating calves. Do not use in calves to be processed for veal. A milk discard time has not been established for use in lactating dairy cattle. Do not use in female dairy cattle 20 months or older, including dry dairy cows. Use in these cattle may cause drug residues in milk and/or in calves born to these cows. Withdraw 5 days before slaughter. Use of more than one product containing neomycin or failure to follow withdrawal times may result in illegal drug residues.		

(1) Mixing directions are for example only and are based on mixing 1 lb of dry milk replacer with 1 gallon of water.

(2) If calf weighs 100 lb, consuming 1 gallon of milk replacer per day.

NADA 138-939

APPROVED BY FDA

FOR USE IN DRY FEEDS ONLY. NOT FOR USE IN LIQUID FEED SUPPLEMENTS.

Store at 25°C (77°F) with excursions permitted to 37°C (98.6°F).

Restricted Drug (California) - Use Only as Directed
Not For Human Use

Distributed by

PharmGate
Animal Health

14040 Industrial Rd. Omaha, NE 68144

